



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OPP OFFICIAL RECORD
HEALTH EFFECTS DIVISION
SCIENTIFIC DATA REVIEWS
EPA SERIES 361

OFFICE OF
PREVENTION, PESTICIDES, AND
TOXIC SUBSTANCES

TXR No. 0054793

MEMORANDUM

DATE: December 12, 2007

SUBJECT: **Cyprosulfamide:** Qualitative Risk Assessment Based On C57BL/6J
Mouse Carcinogenicity Dietary Study

P.C. Code: 877400

TO: John Doherty, Toxicologist
Reregistration Branch 3
Health Effects Division (7509P)

FROM: Lori L. Brunsman, Statistician
Science Information Management Branch
Health Effects Division (7509P)

THROUGH: Jessica Kidwell, EPS
and
Jess Rowland, Branch Chief
Science Information Management Branch
Health Effects Division (7509P)

BACKGROUND

A carcinogenicity study in C57BL/6J mice was conducted by Bayer CropScience, Sophia Antipolis Cedex, France, for Bayer AG, Bayer CropScience, Monheim, Germany, and completed December 1, 2006 (Study No. SA 04065, MRID No. 47069818).

The study design originally allocated groups of 50 mice to dose levels of 0, 350, 2000 and 7000 ppm of Cyprosulfamide for 80 weeks. An additional ten mice per sex per dose were designated for interim sacrifice at week 53. Since there was excessive mortality at the 7000 ppm dose group, an additional dose group of 3500 ppm was added one month after the start of the study and all 7000 ppm dose group animals were terminated (females at week 6 and males at week 46). Final dosing levels were

DEC 20 2007
REC
54

equivalent to 0, 50, 287 and 506 mg/kg/day for males and 0, 63, 354 and 616 mg/kg/day for females. There were no compound-related tumors in the males so only the analyses of the females are presented in this document.

ANALYSES

Survival Analyses

There were no statistically significant incremental changes in mortality with increasing doses of Cyprosulfamide in female mice (Table 1).

Tumor Analyses

Female mice had a statistically significant trend at $p < 0.01$, and a significant pair-wise comparison of the 3500 ppm dose group with the controls at $p < 0.05$, for histiocytic sarcomas. The statistical analyses of the tumors in the female mice were based upon Fisher's Exact Test for pair-wise comparisons and the Exact Test for trend (Table 2).

Table 1. Cyprosulfamide – C57BL/6J Mouse Study (MRID 47069818)

Female Mortality Rates⁺ and Cox or Generalized K/W Test Results

Dose (ppm)	<u>Weeks</u>				Total
	1-26	27-52	53 ⁱ	53-81 ^f	
0	1/60	5/59	10/54	4/44	10/50 (20)
350	2/60	2/58	9/56	4/47	8/51 (16)
2000	2/58 ^a	7/56	7/49	4/42	13/51 (25)
3500	0/60	2/60	10/58	9/48	11/50 (22)

⁺Number of animals that died during interval/Number of animals alive at the beginning of the interval.

ⁱInterim sacrifice at week 53.

^fFinal sacrifice at weeks 79-81.

^aTwo accidental deaths, one each at weeks 2 and 26, dose 2000 ppm.

()Percent.

Note: Time intervals were selected for display purposes only.
 Significance of trend denoted at control.
 Significance of pair-wise comparison with control denoted at dose level.
 If *, then $p < 0.05$. If **, then $p < 0.01$.

Table 2. Cyprosulfamide – C57BL/6J Mouse Study (MRID 47069818)

Female Histiocytic Sarcoma Tumor Rates⁺
and Fisher's Exact Test and Exact Test for Trend Results

	Dose (ppm)			
	0	350	2000	3500
Histiocytic Sarcomas (%)	0/44 (0)	1/47 (2)	0/42 (0)	5 ^a /48 (10)
p =	0.007396**	0.51648	1.0000	0.03482*

+Number of tumor bearing animals/Number of animals examined, excluding those that died or were sacrificed before week 54.

^aFirst histiocytic sarcoma observed at week 62, dose 3500 ppm.

Note: Significance of trend denoted at control.
 Significance of pair-wise comparison with control denoted at dose level.
 If *, then $p < 0.05$. If **, then $p < 0.01$.

References

Cox, D.R. (1972) Regression Models and Life Tables (with discussion). J. Royal Stat. Soc. Ser. B. 34, 187-220.

Gart, J.J., D. Krewski, P.N. Lee, R.E. Tarone, and J. Wahrendorf (1986) The Design and Analysis of Long-Term Animal Experiments. In: Statistical Methods in Cancer Research, Volume III. IARC Scientific Publications No. 79. Lyon, France: International Agency for Research on Cancer, p. 18.

Peto, R., M. Pike, N. Day, R. Gray, P. Lee, S. Parish, J. Peto, S. Richard, and J. Wahrendorf (1980) Guidelines for Simple, Sensitive, Significant Tests for Carcinogenic Effects in Long-Term Animal Experiments. In: Monographs on the long-term and short-term screening assays for carcinogens: a critical appraisal. IARC Monographs, Supplement 2. Lyon, France: International Agency for Research on Cancer, pp. 311-426.

Thomas, D.G., N. Breslow, and J.J. Gart (1977) Trend and Homogeneity Analyses of Proportions and Life Table Data. Computers and Biomedical Research 10, 373-381.



13544



R155880

Chemical: Cyprosulfamide

PC Code:
877400

HED File Code: 31000 Qualitative Analysis Reviews

Memo Date: 12/12/2007

File ID: TX0054793

Accession #: 000-00-0124

HED Records Reference Center
1/15/2008

